

# Nonlocal Cosmology

Ivan Dimitrijević, Branko Dragovich, Jelena Grujić  
and Zoran Rakić

# Nonlocal Modified Gravity

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Nonlocal gravity is a modification of Einstein general relativity in such way that Einstein-Hilbert action contains a function  $f(\square, R)$ . Our action is given by

$$S = \int d^4x \sqrt{-g} \left( \frac{R - 2\Lambda}{16\pi G} + CR^{-1} \mathcal{F}(\square) R \right)$$

where  $\square = \frac{1}{\sqrt{-g}} \partial_\mu \sqrt{-g} g^{\mu\nu} \partial_\nu$ ,  $\mathcal{F}(\square) = \sum_{n=0}^{\infty} f_n \square^n$  and  $C$  is a constant.

We use Friedmann-Lemaître-Robertson-Walker (FLRW) metric

$$ds^2 = -dt^2 + a^2(t) \left( \frac{dr^2}{1-kr^2} + r^2 d\theta^2 + r^2 \sin^2 \theta d\phi^2 \right), \quad k \in \{-1, 0, 1\}.$$

# FLRW metric

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

$$R = \frac{6(a(t)\ddot{a}(t) + \dot{a}(t)^2 + k)}{a(t)^2}$$

In case of FLRW metric the d'Alembert operator is

$$\square R = -\ddot{R} - 3H\dot{R}, \quad H = \frac{\dot{a}}{a}$$

# Nonlocal Model with Term $R^{-1}\mathcal{F}(\square)R$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

We can rewrite our action in the following way

$$S_1 = \int d^4x \sqrt{-g} \left( \frac{R}{16\pi G} + R^{-1} \mathcal{F}(\square) R \right),$$

where  $\mathcal{F}(\square) = \sum_{n=0}^{\infty} f_n \square^n$  and when  $f_0 = -\frac{\Lambda}{8\pi G}$  it plays role of the cosmological constant.

# Nonlocal Model with Term $R^{-1}\mathcal{F}(\square)R$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

We can rewrite our action in the following way

$$S_1 = \int d^4x \sqrt{-g} \left( \frac{R}{16\pi G} + R^{-1}\mathcal{F}(\square)R \right),$$

where  $\mathcal{F}(\square) = \sum_{n=0}^{\infty} f_n \square^n$  and when  $f_0 = -\frac{\Lambda}{8\pi G}$  it plays role of the cosmological constant.

- The nonlocal term  $R^{-1}\mathcal{F}(\square)R$  is invariant under transformation  $R \rightarrow CR$ . It means that effect of nonlocality does not depend on the magnitude of scalar curvature  $R$ , but on its spacetime dependence, and in the FLRW case is sensitive only to dependence of  $R$  on time  $t$ .

# Equations of Motion

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

By variation of action  $S_1$  with respect to metric  $g^{\mu\nu}$  we obtain

$$\begin{aligned} R_{\mu\nu} V - (\nabla_\mu \nabla_\nu - g_{\mu\nu} \square) V - \frac{1}{2} g_{\mu\nu} R^{-1} \mathcal{F}(\square) R \\ + \frac{1}{2} \sum_{n=1}^{\infty} f_n \sum_{l=0}^{n-1} (g_{\mu\nu} (\partial_\alpha \square^l (R^{-1}) \partial^\alpha \square^{n-1-l} R + \square^l (R^{-1}) \square^{n-l} R) \\ - 2 \partial_\mu \square^l (R^{-1}) \partial_\nu \square^{n-1-l} R) = - \frac{G_{\mu\nu}}{16\pi G}, \end{aligned}$$

where  $V = \mathcal{F}(\square) R^{-1} - R^{-2} \mathcal{F}(\square) R$ .

# Equations of Motion

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

The trace and 00-component of EOM are

$$RV + 3\Box V + \sum_{n=1}^{\infty} f_n \sum_{l=0}^{n-1} (\partial_{\alpha} \Box^l (R^{-1}) \partial^{\alpha} \Box^{n-1-l} R + 2\Box^l (R^{-1}) \Box^{n-l} R) - 2R^{-1} \mathcal{F}(\Box) R = \frac{R}{16\pi G},$$

$$R_{00} V - (\nabla_0 \nabla_0 - g_{00} \Box) V - \frac{1}{2} g_{00} R^{-1} \mathcal{F}(\Box) R + \frac{1}{2} \sum_{n=1}^{\infty} f_n \sum_{l=0}^{n-1} (g_{00} (\partial_{\alpha} \Box^l (R^{-1}) \partial^{\alpha} \Box^{n-1-l} R + \Box^l (R^{-1}) \Box^{n-l} R) - 2\partial_0 \Box^l (R^{-1}) \partial_0 \Box^{n-1-l} R) = -\frac{G_{00}}{16\pi G}.$$

# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Let  $R = R_0 = \text{const}$  and we obtain

$$6\left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2}\right) = R_0.$$



# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Let  $R = R_0 = \text{const}$  and we obtain

$$6\left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2}\right) = R_0.$$

The change of variable  $b(t) = a^2(t)$  yields

$$3\ddot{b} - R_0 b = -6k.$$

# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Let  $R = R_0 = \text{const}$  and we obtain

$$6\left(\frac{\ddot{a}}{a} + \left(\frac{\dot{a}}{a}\right)^2 + \frac{k}{a^2}\right) = R_0.$$

The change of variable  $b(t) = a^2(t)$  yields

$$3\ddot{b} - R_0 b = -6k.$$

Depending on the sign of  $R_0$  we have the following solutions for  $b(t)$

$$\begin{aligned} R_0 > 0 \quad b(t) &= \frac{6k}{R_0} + \sigma e^{\sqrt{\frac{R_0}{3}}t} + \tau e^{-\sqrt{\frac{R_0}{3}}t} \\ R_0 < 0 \quad b(t) &= \frac{6k}{R_0} + \sigma \cos \sqrt{\frac{-R_0}{3}}t + \tau \sin \sqrt{\frac{-R_0}{3}}t \end{aligned}$$

# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

When we set  $R = R_0 = \text{const}$  into trace and 00-equation we obtain the following system

$$-2f_0 = \frac{R_0}{16\pi G}, \quad \frac{1}{2}f_0 = -\frac{G_{00}}{16\pi G}$$

# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

When we set  $R = R_0 = \text{const}$  into trace and 00-equation we obtain the following system

$$-2f_0 = \frac{R_0}{16\pi G}, \quad \frac{1}{2}f_0 = -\frac{G_{00}}{16\pi G}$$

The last system has a solution iff

$$R_0 + 4R_{00} = 0.$$

Note that  $R_{00}$  can be written in terms of function  $b(t)$  as

$$R_{00} = -\frac{3\ddot{a}}{a} = \frac{3((\dot{b})^2 - 2b\ddot{b})}{4b^2}.$$

# Cosmological solutions with constant scalar curvature

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Now, from  $R_0 + 4R_{00} = 0$  we obtain the following conditions on the parameters  $\sigma$  and  $\tau$ :

$$\begin{aligned} R_0 > 0 & \quad 9k^2 = R_0^2 \sigma \tau, \\ R_0 < 0 & \quad 36k^2 = R_0^2 (\sigma^2 + \tau^2). \end{aligned}$$

## Case 1: $R_0 < 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

If  $k = -1$  we can define  $\varphi$  by  $\sigma = \frac{-6}{R_0} \cos \varphi$  and  $\tau = \frac{-6}{R_0} \sin \varphi$  and rewrite  $a(t)$  and  $b(t)$  as

$$b(t) = \frac{-12}{R_0} \cos^2 \frac{1}{2} \left( \sqrt{-\frac{R_0}{3}} t - \varphi \right),$$
$$a(t) = \sqrt{\frac{-12}{R_0}} \left| \cos \frac{1}{2} \left( \sqrt{-\frac{R_0}{3}} t - \varphi \right) \right|.$$

## Case 1: $R_0 < 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

If  $k = -1$  we can define  $\varphi$  by  $\sigma = \frac{-6}{R_0} \cos \varphi$  and  $\tau = \frac{-6}{R_0} \sin \varphi$  and rewrite  $a(t)$  and  $b(t)$  as

$$b(t) = \frac{-12}{R_0} \cos^2 \frac{1}{2} \left( \sqrt{-\frac{R_0}{3}} t - \varphi \right),$$
$$a(t) = \sqrt{\frac{-12}{R_0}} \left| \cos \frac{1}{2} \left( \sqrt{-\frac{R_0}{3}} t - \varphi \right) \right|.$$

In the last case  $k = +1$  we can transform  $b(t)$  to

$$b(t) = \frac{12}{R_0} \sin^2 \frac{1}{2} \left( \sqrt{-\frac{R_0}{3}} t - \varphi \right),$$

which is non positive and hence yields no solutions.

## Case 2: $R_0 > 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Set  $k = 0$  then we obtain a solution with constant Hubble parameter. Alternatively, if we set  $k = +1$  we can find  $\varphi$  such that  $\sigma + \tau = \frac{6}{R_0} \cosh \varphi$  and  $\sigma - \tau = \frac{6}{R_0} \sinh \varphi$ . Moreover, we obtain

$$b(t) = \frac{12}{R_0} \cosh^2 \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right),$$
$$a(t) = \sqrt{\frac{12}{R_0}} \cosh \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right).$$



## Case 2: $R_0 > 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Set  $k = 0$  then we obtain a solution with constant Hubble parameter. Alternatively, if we set  $k = +1$  we can find  $\varphi$  such that  $\sigma + \tau = \frac{6}{R_0} \cosh \varphi$  and  $\sigma - \tau = \frac{6}{R_0} \sinh \varphi$ . Moreover, we obtain

$$b(t) = \frac{12}{R_0} \cosh^2 \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right),$$
$$a(t) = \sqrt{\frac{12}{R_0}} \cosh \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right).$$

At the end if we set  $k = -1$  we can transform  $b(t)$  to

$$b(t) = \frac{12}{R_0} \sinh^2 \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right),$$
$$a(t) = \sqrt{\frac{12}{R_0}} \left| \sinh \frac{1}{2} \left( \sqrt{\frac{R_0}{3}} t + \varphi \right) \right|.$$

## Case 2.1: $R = 12\lambda^2$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

We consider the scale factor of the form

$$a(t) = a_0(\sigma_1 e^{\lambda t} + \tau_1 e^{-\lambda t}).$$

## Case 2.1: $R = 12\lambda^2$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

We consider the scale factor of the form

$$a(t) = a_0(\sigma_1 e^{\lambda t} + \tau_1 e^{-\lambda t}).$$

We have

$$H(t) = \frac{\lambda(e^{2\lambda t}\sigma_1 - \tau_1)}{e^{2\lambda t}\sigma_1 + \tau_1},$$
$$R(t) = \frac{6(e^{2\lambda t}k + 2\lambda^2(e^{4\lambda t}\sigma_1^2 + \tau_1^2)a_0^2)}{(e^{2\lambda t}\sigma_1 + \tau_1)^2 a_0^2}.$$

In order to have  $R = \text{const}$  we have to satisfy condition  
 $k = 4\lambda^2 a_0^2 \sigma_1 \tau_1$ .

From the last condition we obtain  $R = 12\lambda^2$ .

## Case 2.1: $R = 12\lambda^2$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Substituting this into equations trace and 00-component we obtain

$$f_0 = -\frac{3\lambda^2}{8\pi G}, \quad f_i \in \mathbb{R}, \quad i \geq 1.$$

## Case 2.1: $R = 12\lambda^2$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Substituting this into equations trace and 00-component we obtain

$$f_0 = -\frac{3\lambda^2}{8\pi G}, \quad f_i \in \mathbb{R}, \quad i \geq 1.$$

In particular, if we set  $\sigma_1 = \tau_1 = \frac{1}{2}$  the scale factor becomes

$$a(t) = a_0 \cosh(\lambda t).$$

In this case from condition  $k = 4\lambda^2 a_0^2 \sigma_1 \tau_1$  we see that the only nontrivial case is when  $k$  is equal to 1. From this we obtain  $a_0 = \frac{1}{\lambda}$ .

## Case 2.1: $R = 12\lambda^2$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Substituting this into equations trace and 00-component we obtain

$$f_0 = -\frac{3\lambda^2}{8\pi G}, \quad f_i \in \mathbb{R}, \quad i \geq 1.$$

In particular, if we set  $\sigma_1 = \tau_1 = \frac{1}{2}$  the scale factor becomes

$$a(t) = a_0 \cosh(\lambda t).$$

In this case from condition  $k = 4\lambda^2 a_0^2 \sigma_1 \tau_1$  we see that the only nontrivial case is when  $k$  is equal to 1. From this we obtain  $a_0 = \frac{1}{\lambda}$ . If we take  $\sigma_1 = 0$  or  $\tau_1 = 0$  the scale factor becomes

$$a(t) = a_0 e^{\lambda t}.$$

From condition  $k = 4\lambda^2 a_0^2 \sigma_1 \tau_1$  we see that in this case  $k$  must be equal to 0.

## Case 3: $R_0 = 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

- The case  $R_0 = 0$  can be considered as limit of  $R_0 \rightarrow 0$  in both cases  $R_0 < 0$  and  $R_0 > 0$ .

## Case 3: $R_0 = 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

- The case  $R_0 = 0$  can be considered as limit of  $R_0 \rightarrow 0$  in both cases  $R_0 < 0$  and  $R_0 > 0$ .
- When  $R_0 < 0$  there is condition  $36k^2 = R_0^2(\sigma^2 + \tau^2)$ . From this condition,  $R_0 \rightarrow 0$  implies  $k = 0$  and arbitrary values of constants  $\sigma$  and  $\tau$ . The same conclusion obtains when  $R_0 > 0$  with condition  $9k^2 = R_0^2\sigma\tau$ . In both these cases there is Minkowski solution with  $b(t) = \text{constant} > 0$  and consequently  $a(t) = \text{constant} > 0$ .



## Case 3: $R_0 = 0$

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

- The case  $R_0 = 0$  can be considered as limit of  $R_0 \rightarrow 0$  in both cases  $R_0 < 0$  and  $R_0 > 0$ .
- When  $R_0 < 0$  there is condition  $36k^2 = R_0^2(\sigma^2 + \tau^2)$ . From this condition,  $R_0 \rightarrow 0$  implies  $k = 0$  and arbitrary values of constants  $\sigma$  and  $\tau$ . The same conclusion obtains when  $R_0 > 0$  with condition  $9k^2 = R_0^2\sigma\tau$ . In both these cases there is Minkowski solution with  $b(t) = \text{constant} > 0$  and consequently  $a(t) = \text{constant} > 0$ .
- Note that the Minkowski space solution can be also obtained from the case  $R = 12\lambda^2$ . Namely, the solution  $a(t) = a_0 e^{\lambda t}$  satisfies  $H = \lambda$ . Taking the limit  $\lambda \rightarrow 0$  in  $a(t) = a_0 e^{\lambda t}$  one obtains Minkowski space as a solution for

$$f_0 = 0, \quad f_i \in \mathbb{R}, \quad i \geq 1.$$

# Concluding remarks

Nonlocal  
Cosmology







Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

- We presented cosmological solutions for constant scalar curvature of model given by

$$S_1 = \int d^4x \sqrt{-g} \left( \frac{R}{16\pi G} + R^{-1} \mathcal{F}(\square) R \right).$$

- When  $R = R_0 < 0$  there is nontrivial solution  $a(t) = \sqrt{\frac{-12}{R_0}} |\cos \frac{1}{2}(\sqrt{-\frac{R_0}{3}} t - \varphi)|$  for  $k = -1$ .
- In the case  $R = R_0 > 0$  there are solutions for all three values of curvature constant  $k = 0, \pm 1$ .
- The case  $R = R_0 = 0$  was considered as limit of  $R_0 \rightarrow 0$  in both cases  $R_0 < 0$  and  $R_0 > 0$ , and Minkowski space solution was obtained.
- All obtained solutions are defined for all values of cosmic time  $t$ .
- Solutions for  $R_0 > 0$  with  $k = 0, +1$  are nonsingular bounce cosmological solutions.
- Solution  $a(t) = \sqrt{\frac{-12}{R_0}} |\cos \frac{1}{2}(\sqrt{-\frac{R_0}{3}} t - \varphi)|$ , which is for  $R_0 < 0$  and  $k = -1$ , is a singular cyclic solution.

# References

-  Biswas, T., Mazumdar, A., Siegel, W: Bouncing universes in string-inspired gravity. J. Cosmology Astropart. Phys. 0603, 009 (2006) [arXiv:hep-th/0508194]
-  Biswas, T., Koivisto, T., Mazumdar, A.: Towards a resolution of the cosmological singularity in non-local higher derivative theories of gravity. J. Cosmology Astropart. Phys. 1011, 008 (2010) [arXiv:1005.0590v2 [hep-th]]
-  Koshelev, A.S., Vernov, S.Yu.: On bouncing solutions in non-local gravity. [arXiv:1202.1289v1 [hep-th]]
-  Dimitrijevic, I., Dragovich, B., Grujic J., Rakić, Z.: New cosmological solutions in nonlocal modified gravity. Rom. Journ. Phys. 58 (5-6), 550559 (2013) [arXiv:1302.2794 [gr-qc]]
-  T. Biswas, A. S. Koshelev, A. Mazumdar, S. Yu. Vernov, “Stable bounce and inflation in non-local higher derivative cosmology”, JCAP **08** (2012) 024, [arXiv:1206.6374v2 [astro-ph.CO]].
-  I. Dimitrijevic, B. Dragovich, J. Grujic, Z. Rakić, “A new model of nonlocal modified gravity”, Publications de l’Institut Mathematique **94** (108), 187–196 (2013).

# Motivation

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

Large cosmological observational findings:

- High orbital speeds of galaxies in clusters. (F.Zwicky, 1933)
- High orbital speeds of stars in spiral galaxies. (Vera Rubin, at the end of 1960es)
- Accelerated expansion of the Universe. (1998)

# Problem solving approaches

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

There are two problem solving approaches:

- Dark matter and energy
- Modification of Einstein theory of gravity

$$R_{\mu\nu} - \frac{1}{2}Rg_{\mu\nu} = 8\pi GT_{\mu\nu} - \Lambda g_{\mu\nu}, c = 1$$

where  $T_{\mu\nu}$  is stress-energy tensor,  $g_{\mu\nu}$  are the elements of the metric tensor,  $R_{\mu\nu}$  is Ricci tensor and  $R$  is scalar curvature of metric.

# Dark matter and energy

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

- If Einstein theory of gravity can be applied to the whole Universe then the Universe contains about 5% of ordinary matter, 27% of dark matter and 68% of dark energy.
- It means that 95% of total matter, or energy, represents dark side of the Universe, which nature is unknown.
- Dark matter is responsible for orbital speeds in galaxies, and dark energy is responsible for accelerated expansion of the Universe.

# Modification of Einstein theory of gravity

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

## Motivation for modification of Einstein theory of gravity

- The validity of General Relativity on cosmological scale is not confirmed.
- Dark matter and dark energy are not yet detected in the laboratory experiments.
- Another cosmological problem is related to the Big Bang singularity. Namely, under rather general conditions, general relativity yields cosmological solutions with zero size of the universe at its beginning, what means an infinite matter density.
- Note that when physical theory contains singularity, it is not valid in the vicinity of singularity and must be appropriately modified.

# Approaches to modification of Einstein theory of gravity

Nonlocal  
Cosmology

Ivan Dimitrijević,  
Branko  
Dragovich,  
Jelena Grujić  
and Zoran Rakić

There are different approaches to modification of Einstein theory of gravity.

- Einstein General Theory of Relativity

From action  $S = \int \left( \frac{R}{16\pi G} - L_m - 2\Lambda \right) \sqrt{-g} d^4x$  using variational methods we get field equations

$$R_{\mu\nu} - \frac{1}{2} R g_{\mu\nu} = 8\pi G T_{\mu\nu} - \Lambda g_{\mu\nu}, \quad c = 1$$

where  $T_{\mu\nu}$  is stress-energy tensor,  $g_{\mu\nu}$  are the elements of the metric tensor,  $R_{\mu\nu}$  is Ricci tensor and  $R$  is scalar curvature of metric.

Currently there are mainly two approaches:

- f(R) Modified Gravity
- Nonlocal Gravity